## ROUTINE ASSESSMENT OF PATIENT INDEX DATA

The RAPID3 includes a subset of core variables found in the Multi-dimensional HAQ (MD-HAQ). Page 1 of the MD-HAQ, shown here, includes an assessment of physical function (section 1), a patient global assessment (PGA) for pain (section 2), and a PGA for global health (section 3). RAPID3 scores are quickly tallied by adding subsets of the MD-HAQ as follows:

1. please check the <b>ONE</b> best answer for your abilities at this time:							
OVER THE LAST WEEK, WERE YOU ABLE TO:	WITHOUT <b>ANY</b> DIFFICULTY	WITH <b>SOME</b> DIFFICULTY	WITH <b>MUCH</b> DIFFICULTY	<b>UNABLE</b> TO DO			
a. Dress yourself, including tying shoelaces and doing buttons?	0	1	2	3			
<b>b.</b> Get in and out of bed?	0	1	2	3			
c. Lift a full cup or glass to your mouth?	0	1	2	3			
d. Walk outdoors on flat ground?	0	1	2	3			
e. Wash and dry your entire body?	0	1	2	3			
f. Bend down to pick up clothing from the floor?	0	1	2	3			
g. Turn regular faucets on and off?	0	1	2	3			
h. Get in and out of a car, bus, train, or airplane?	0	1	2	3			
i. Walk two miles or three kilometers, if you wish?	0	1	2	3			
j. Participate in recreational activities and sports as you would like, if you wish?	0	1	2	3			
k. Get a good night's sleep?	0	1.1	2.2	3.3			
1. Deal with feelings of anxiety or being nervous?	0	1.1	2.2	3.3			
m. Deal with feelings of depression or feeling blue?	0	1.1	2.2	3.3			

1 asi FN	N (0-10):
1. a-, 11	<b>1</b> (0-10).
1=0.3	16=5.3
2=0.7	
3=1.0	18=6.0
4=1.3	19=6.3
5=1.7	20=6.7
6=2.0	21=7.0
7=2.3	22=7.3
8=2.7	23=7.7
9=3.0	24=8.0
10=3.3	25=8.3
11=3.7	26=8.7
12=4.0	27=9.0
13=4.3	28=9.3
14=4.7	29=9.7
15=5.0	30=10
2. PN (	0-10).
	E (0-10):
3. PTG	
3. PTG	
3. PIG	
	3 (0-30)

## 2. HOW MUCH PAIN HAVE YOU HAD BECAUSE OF YOUR CONDITION **OVER THE PAST WEEK?**Please indicate below how severe your pain has been:

NO	PAIN														PA	IN AS	BAD A	AS IT C	COULI	) BE
0	0.5	1.0	1.5	2.0	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	6.5	<b>7.0</b>	7.5	8.0	8.5	9.0	9.5	10

# 3. Considering all the ways in which illness and health conditions may affect you at this time, please indicate below how you are doing:

VER	Y WEL	L																VER	Y POO	RLY
0	0.5	1.0	1.5	2.0	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	6.5	<b>7.0</b>	7.5	8.0	8.5	9.0	9.5	10

CONVERSION TABLE

Near Remission (NR): 1=0.3; 2=0.7; 3=1.0 Low Severity (LS): 4=1.3; 5=1.7; 6=2.0 Moderate Severity (MS): 7=2.3; 8=2.7; 9=3.0

Moderate Severity (MS): 7=2.3; 8=2.7; 9=3.0; 10=3.3; 11=3.7; 12=4.0

High Severity (HS): 13-4.3; 14-4.7; 15-5.0; 16-5.3; 17-5.7; 18-6.0; 19-6.3; 20-6.7; 21-7.0; 22-7.3; 23-7.7; 24-8.0; 25-8.3; 26-8.7; 27-9.0; 28-9.3; 29-9.7; 30-10.0

### HOW TO CALCULATE RAPID 3 SCORES

- 1. Ask the patient to complete questions 1, 2, and 3 while in the waiting room prior to his/her visit.
- 2. For question 1, add up the scores in questions A-J only (questions K-M have been found to be informative, but are not scored formally). Use the formula in the box on the right to calculate the formal score (0-10). For example, a patient whose answers total 19 would score a 6.3. Enter this score as an evaluation of the patient's functional status (FN).
- 3. For question 2, enter the raw score (0-10) in the box on the right as an evaluation of the patient's pain tolerance (PN).
- 4. For question 3, enter the raw score (0-10) in the box on the right as an evaluation of the patient's global estimate (PTGE).
- **5.** Add the total score (0-30) from questions 1, 2, and 3 and enter them as the patient's RAPID 3 cumulative score. Use the final conversion table to simplify the patient's weighed RAPID 3 score. For example, a patient who scores 11 on the cumulative RAPID 3 scale would score a weighed 3.7. A patient who scores between 0–1.0 is defined as near remission (NR); 1.3–2.0 as low severity (LS); 2.3–4.0 as moderate severity (MS); and 4.3–10.0 as high severity (HS).


# ROUTINE ASSESSMENT OF PATIENT INDEX DAT

The RAPID includes a subset of core variables found in the Multi-dimensional HAQ (MD-HAQ). Page 1 of the MD-HAQ, shown here, includes an assessment of physical function (section 1), a patient global assessment (PGA) for pain (section 2), and a PGA for global health (section 3). RAPID scores are quickly tallied by adding subsets of the MD-HAQ as follows:


1. please check the <b>ONE</b> best ans	SWER FOR YO	UR ABILITIES	S AT THIS TIM	ιE:
OVER THE LAST WEEK, WERE YOU ABLE TO:	WITHOUT <b>ANY</b> DIFFICULTY	WITH <b>SOME</b> DIFFICULTY	WITH <b>MUCH</b> DIFFICULTY	<b>UNABLE</b> TO DO
a. Dress yourself, including tying shoelaces and doing buttons?	<u>×</u> 0	1	2	3
b. Get in and out of bed?	0	<u>×</u> 1	2	3
c. Lift a full cup or glass to your mouth?	0	1	2	×3
d. Walk outdoors on flat ground?	<u>×</u> 0	1	2	3
e. Wash and dry your entire body?	<u>×</u> 0	1	2	3
f. Bend down to pick up clothing from the floor?	0	<u>×</u> 1	2	3
g. Turn regular faucets on and off?	0	1	<u>×</u> 2	3
h. Get in and out of a car, bus, train, or airplane?	0	1	<u>×</u> 2	3
i. Walk two miles or three kilometers, if you wish?	0	<u>×</u> 1	2	3
j. Participate in recreational activities and sports as you would like, if you wish?	0	<u>×_1</u>	2	3
k. Get a good night's sleep?	<u>×</u> 0	1.1	2.2	3.3
1. Deal with feelings of anxiety or being nervous?	0	1.1	2.2	× 3.3
m. Deal with feelings of depression or feeling blue?	0	1.1	<u>×</u> 2.2	3.3

,	N (0-10):
3.7	7
5=1.7 6=2.0 7=2.3 8=2.7 9=3.0 10=3.3 11=3.7 12=4.0 13=4.3 14=4.7	17=5.7 18=6.0 19=6.3 20=6.7 21=7.0 22=7.3 23=7.7 24=8.0 25=8.3 26=8.7 27=9.0 28=9.3 29=9.7
15=5.0 2. PN ((	0-10):
3. PTG	E (0-10):
1.0	)
RAPID	3 (0-30) . <b>2</b>

### 2. HOW MUCH PAIN HAVE YOU HAD BECAUSE OF YOUR CONDITION OVER THE PAST WEEK? Please indicate below how severe your pain has been:



### 3. considering all the ways in which illness and health conditions may affect you AT THIS TIME, PLEASE INDICATE BELOW HOW YOU ARE DOING:



**CONVERSION TABLE** 

Near Remission (NR): 1=0.3; 2=0.7; 3=1.0 Low Severity (LS): 4=1.3; 5=1.7; 6=2.0

Moderate Severity (MS): 7=2.3; 8=2.7; 9=3.0; 10=3.3; 11=3.7; 12=4.0

High Severity (HS): 13=4.3; 14=4.7; 15=5.0; 16=5.3; 17=5.7; 18=6.0; 19=6.3; 20=6.7; 21=7.0; 22=7.3; 23=7.7; 24=8.0; 25=8.3; 26=8.7; 27=9.0; 28=9.3; 29=9.7; 30=10.0

### HOW TO CALCULATE RAPID 3 SCORES

- 1. Ask the patient to complete questions 1, 2, and 3 while in the waiting room prior to his/her visit.
- 2. For question 1, add up the scores in questions A-J only (questions K-M have been found to be informative, but are not scored formally). Use the formula in the box on the right to calculate the formal score (0-10). For example, a patient whose answers total 19 would score a 6.3. Enter this score as an evaluation of the patient's functional status (FN).
- 3. For question 2, enter the raw score (0-10) in the box on the right as an evaluation of the patient's pain tolerance (PN).
- 4. For question 3, enter the raw score (0-10) in the box on the right as an evaluation of the patient's global estimate (PTGE).
- 5. Add the total score (0-30) from questions 1, 2, and 3 and enter them as the patient's RAPID 3 cumulative score. Use the final conversion table to simplify the patient's weighed RAPID 3 score. For example, a patient who scores 11 on the cumulative RAPID 3 scale would score a weighed 3.7. A patient who scores between 0-1.0 is defined as near remission (NR); 1.3-2.0 as low severity (LS); 2.3-4.0 as moderate severity (MS); and 4.3-10.0 as high severity (HS).